

Experimental Physiology

<https://ep.msubmit.net>

EP-SC-2019-088087R1

Title: The Effects of Long-Term Muscle Disuse on Neuromuscular Function in Unilateral Transtibial Amputees

Authors: Amy Rebecca Sibley
Siobhan Strike
Sarah Catherine Moudy
Neale Anthony Tillin

Author Conflict: No competing interests declared

Running Title: Long-Term Muscular Disuse in Unilateral Transtibial Amputees

Abstract: The purpose of this study was to determine: (1) whether individuals with unilateral transtibial amputations (ITTAs), who habitually disuse the quadriceps muscles of their amputated limb, provide an effective model for assessing the effects of long-term muscle disuse; and (2) the effects of such disuse on quadriceps muscle strength and neuromuscular function in this population. Nine ITTAs and nine controls performed isometric voluntary knee extensions in both limbs to assess maximal voluntary torque (MVT) and rate of torque development (RTD). The interpolated twitch technique and EMG normalised to maximal M-wave assessed neural activation, involuntary (twitch and octet) contractions assessed intrinsic contractile properties, and ultrasound images of the vastus lateralis assessed muscle architecture. Clinical gait analysis was used to measure knee kinetic data during walking at an habitual speed. ITTAs displayed 54-60% lower peak knee extensor moments during walking in the amputated than intact/control

limbs, but the intact and control limbs were comparable for loading during walking and muscle strength variables, suggesting the intact limb provides a suitable internal control for comparison to the disused amputated limb. MVT and RTD were ~60% and ~75% lower, respectively, in the amputated than intact/control limbs. The differences in MVT appeared associated with ~40% and ~43% lower muscle thickness and neural activation, respectively, whilst the differences in RTD appeared associated with the decline in MVT coupled with slowing of the intrinsic contractile properties. These results indicate considerable changes in strength and neuromuscular function with long-term disuse, that could not be predicted from short-term disuse studies.

New Findings: What is the central question of this study? The effects of long-term muscle disuse on neuromuscular function are unclear because disuse studies are typically short-term. This study used a novel model (unilateral transtibial amputees) to investigate the effects of long-term disuse on quadriceps neuromuscular function. What is the main finding and its importance? Kinetic analysis (knee extension moments during gait) indicated habitual disuse of the amputated limb quadriceps, accompanied by lower quadriceps muscle strength (60-76%) and neural activation (32-44%), slower contractile properties, and altered muscle architecture in the amputated limb, which could not be predicted from short-term disuse studies.

Dual Publication: No

Funding: University of Roehampton: Amy Rebecca Sibley, n/a This study was supported by funds from a PhD scholarship (University of Roehampton Vice-Chancellor's Doctoral Scholarship).

The Effects of Long-Term Muscle Disuse on Neuromuscular Function in Unilateral Transtibial Amputees

Amy R. Sibley,^{1,2} Siobhán Strike,¹ Sarah C. Moudy,^{1,3} Neale A. Tillin¹

¹Department of Life Sciences, University of Roehampton, London, UK

²Department of Allied Health Sciences, London South Bank University, London, UK

³Department of Family Medicine, University of North Texas Health Science Center, Fort
Worth, Texas, USA

Correspondence Address:

Amy R. Sibley

Department of Allied Health Sciences

103 Borough Road

London, UK SE1 0AA

Email: amy.sibley@lsbu.ac.uk

Key words: muscle strength, muscle atrophy, neural activation

Word count: 7596

Number of references: 53

Subject area: Muscle physiology

21 *What is the central question of this study?*

22 The effects of long-term muscle disuse on neuromuscular function are unclear because
23 disuse studies are typically short-term. This study used a novel model (unilateral transtibial
24 amputees) to investigate the effects of long-term disuse on quadriceps neuromuscular
25 function.

26 *What is the main finding and its importance?*

27 Kinetic analysis (knee extension moments during gait) indicated habitual disuse of the
28 amputated limb quadriceps, accompanied by lower quadriceps muscle strength (60-76%)
29 and neural activation (32-44%), slower contractile properties, and altered muscle
30 architecture in the amputated limb, which could not be predicted from short-term disuse
31 studies.

Abstract

The purpose of this study was to determine: (1) whether individuals with unilateral transtibial amputations (ITTAs), who habitually disuse the quadriceps muscles of their amputated limb, provide an effective model for assessing the effects of long-term muscle disuse; and (2) the effects of such disuse on quadriceps muscle strength and neuromuscular function in this population. Nine ITTAs and nine controls performed isometric voluntary knee extensions in both limbs to assess maximal voluntary torque (MVT) and rate of torque development (RTD). The interpolated twitch technique and EMG normalised to maximal M-wave assessed neural activation, involuntary (twitch and octet) contractions assessed intrinsic contractile properties, and ultrasound images of the vastus lateralis assessed muscle architecture. Clinical gait analysis was used to measure knee kinetic data during walking at an habitual speed. ITTAs displayed 54-60% lower peak knee extensor moments during walking in the amputated than intact/control limbs, but the intact and control limbs were comparable for loading during walking and muscle strength variables, suggesting the intact limb provides a suitable internal control for comparison to the disused amputated limb. MVT and RTD were ~60% and ~75% lower, respectively, in the amputated than intact/control limbs. The differences in MVT appeared associated with ~40% and ~43% lower muscle thickness and neural activation, respectively, whilst the differences in RTD appeared associated with the decline in MVT coupled with slowing of the intrinsic contractile properties. These results indicate considerable changes in strength and neuromuscular function with long-term disuse, that could not be predicted from short-term disuse studies.

53 **Introduction**

54 Prolonged disuse of skeletal muscle poses a considerable threat to neuromuscular
55 functional capacity and health (Narici & de Boer 2011). Just nine days of disuse causes
56 considerable reductions in muscle strength, typically measured as maximum voluntary
57 torque (MVT; Rozier et al. 1979) or rate of torque development during contractions
58 performed from rest (RTD; Bamman et al. 1998). The knee extensor (quadriceps) muscles
59 are particularly susceptible to degenerative changes resulting from disuse owing to their
60 large contributions to locomotion, and so are frequently investigated in typical study models
61 of disuse including spaceflight, unilateral lower-limb suspension (ULLS), limb immobilisation,
62 bed rest, and immobilisation during intensive care following surgery (Narici & de Boer 2011).
63 Studies show reductions in quadriceps MVT of approximately 2% per day for the first ten
64 days (Berg & Tesch 1996, Puthuchear et al. 2017, Rozier et al. 1979), slowing to ~1% per
65 week for up to 30 days, with an eventual plateau resulting in average strength losses of
66 around 23% after 120 days of disuse (Dirks et al. 2016, Horstman et al. 2012, Narici & de
67 Boer 2011). The effects of disuse on RTD have not been widely studied, yet RTD may be a
68 more functionally relevant than MVT during rapid human movements such as recovering
69 from a trip or loss of balance (Pijnappels et al. 2008, Behan, Pain & Folland 2018). Long-
70 term muscle disuse is a default position for many clinical populations (Brown et al. 2004) and
71 the sedentary, yet it is unclear how both MVT and RTD change with long-term, habitual
72 disuse, as typical disuse study models last <90-120 days for logistical and ethical reasons.

73 Individuals with unilateral transtibial amputations (ITTAs, below-knee amputation on one
74 limb) may provide a useful model for studying the effects of long-term, habitual disuse.
75 ITTAs adopt an asymmetrical loading pattern characterised by considerably lower vertical
76 ground reaction forces (vGRF) and knee extensor moments on the amputated compared to
77 the intact limb, during movements such as walking (Fey & Neptune 2012), jumping
78 (Schoeman, Diss & Strike 2012), and stair ascent/descent (Schmalz, Blumentritt & Marx
79 2007). This suggests the quadriceps of the amputated limb in ITTAs are chronically disused,

which would explain observations of considerably lower (~50%) quadriceps MVT (Isakov et al. 1996, Lloyd et al. 2010, Pedrinelli et al. 2002) and size (Moirenfeld et al. 2000) in the amputated, compared to the intact and control limbs. Comparison of quadriceps neuromuscular function in the amputated vs. intact limb of ITTAs, coupled with comparison of limb loading during gait as an estimation of typical use, may therefore offer new insight into the long-term effects of habitual disuse. However, currently it is unclear whether the intact limb provides an internal control that is unaffected by the amputation and comparable to the limb of an able-bodied control, which would support the efficacy of ITTAs as a study model of long-term disuse. Despite similar peak vGRF and knee moments during walking gait (Lloyd et al. 2010, Nolan et al. 2003, Sanderson & Martin 1997), previous studies in ITTAs have shown lower MVT in the intact limb compared to able-bodied participant limbs (Isakov et al. 1996, Lloyd et al. 2010, Pedrinelli et al. 2002, Powers et al. 1996). However, the latter studies did not control for other factors known to independently affect muscle strength between the groups such as ageing, health, and sedentary lifestyle (Narici & de Boer 2011, Sacchetti et al. 2013).

Six studies (Isakov et al. 1996, Lloyd et al. 2010, Moirenfeld et al. 2000, Pedrinelli et al. 2002, Powers et al. 1996, Renstrom et al. 1983) have previously measured quadriceps MVT in ITTAs, and none have assessed the changes in RTD in this population. Furthermore, the neuromuscular mechanisms of the considerable strength loss in the amputated limb of ITTAs have not been investigated. Neural activation, assessed via electromyography (EMG) amplitude or the interpolated twitch technique, is considered an important determinant of both MVT and RTD (Balshaw et al. 2016, Folland, Buckthorpe & Hannah 2014, Tillin, Pain & Folland 2011). However, evidence for changes in neural drive with short-term (≤ 89 days) disuse are equivocal with some studies reporting a decrease (Narici & de Boer 2011, Lambertz et al. 2001) and others no change (de Boer et al. 2007, Campbell et al. 2013). RTD also appears to be determined by the intrinsic contractile-speed properties of the muscle, such as RTD relative to peak torque recorded during electrically evoked-involuntary

contractions (e.g. twitch or octets; Folland et al., 2014), and short-term disuse causes a shift towards faster contractile properties (Lambertz et al. 2001). Finally, the maximum force generating potential of a muscle is dependent upon its architecture (Blazevich et al. 2009), and just 21-30 days of disuse have elicited changes such as declines in muscle size ($\leq 10\%$), pennation angle ($\leq 13\%$), and fascicle length ($\leq 9\%$; de Boer et al. 2007, Campbell et al. 2013). Determining the degree of change in these neuromuscular determinants of muscle strength with long-term habitual disuse may allow better targeting of preventative and rehabilitative interventions for populations subject to muscle disuse.

The first aim of this study was to assess the efficacy of unilateral ITTAs as a model to study long-term habitual disuse, by comparing knee-extensor strength (MVT and RTD) and loading (knee extensor moments and impulse) during walking gait of the intact limb with a control able-bodied population, where both groups are healthy, young, and active. The second aim was to assess MVT and RTD, and the neuromuscular determinants of these (neural drive; intrinsic contractile properties; and vastus lateralis muscle architecture) in the disused quadriceps muscles of ITTAs, in comparison to both the intact and an able-bodied control limb.

Methods

Ethical Approval

Participants provided written informed consent prior to their involvement in the study, which complied with the standards set by the 2013 Declaration of Helsinki (except for registration in a database) and was approved by the University of Roehampton Ethics Committee (LSC 16/176) and the NHS Health Research Authority (17/NW/0566).

Participants

Nine male ITTAs and nine male controls took part in this study. Prior to data analysis, groups were matched to ensure similar group means and variability in age, height, body

mass, and physical activity. Physical activity was assessed using the International Physical Activity Questionnaire (Short Format, <http://ipaq.ki.se/downloads.htm>). ITTAs were included if they had a unilateral transtibial amputation performed >6 months prior to involvement in the study, to ensure established ambulation and long-term disuse in the residual limb. ITTAs were excluded if they experienced any discomfort in the residual limb whilst using their prosthesis, and/ or if their amputation occurred due to congenital disorders, or complications arising from metabolic or vascular conditions (e.g. diabetes). Exclusion criteria for both groups included cardiovascular disease risk factors or neuro-musculoskeletal injuries (other than a transtibial amputation in the ITTAs).

Overview

Participants visited the laboratory on three separate occasions, with each visit 3-7 days apart, to complete a familiarisation (visit 1; identical to visit 2), neuromuscular function assessment of the quadriceps muscles of both limbs (visit 2), and a gait assessment (visit 3). Limb order for neuromuscular assessment was randomised. All three sessions commenced at a consistent time (± 2 hours) of day for each participant, following at least 36 hours without strenuous exercise, and 24 hours without alcohol.

Experimental Setup and Measurements

Knee Extension Torque

Isometric strength data were collected using an isokinetic dynamometer (Humac Norm, Computer Sports Medicine Inc., Massachusetts, USA). Participants were seated with a hip angle of 100° (full extension = 180°) and with adjustable straps across the pelvis and shoulders tightened to ensure no extraneous movement. The knee joint angle was set so that the angle during active extension was 110° . Some basic modifications were made to minimise knee joint angle changes during isometric contractions, including the use of a dense foam padding on the seat and limb attachment, and a custom-made lower limb

attachment which could be tightly clamped to the crank arm to remove unnecessary rotation around the crank arm. In all participants, the limb attachment was placed as distal on the tibia as anatomy and participant comfort permitted. For the amputated limb, the crank arm was flipped by 180° to account for the shorter residual tibia.

The analogue torque signal was sampled at 2000 Hz using an external A/D converter (16-bit signal recording resolution; Micro 1401, CED, Cambridge, UK) and interfaced with a PC using Spike 2 software (version 8; CED). Off-line, torque was filtered using a fourth-order low pass Butterworth filter with a cut-off frequency of 10 Hz and corrected for the passive weight of the limb.

Electromyography (EMG)

Electromyography signals were recorded from the superficial knee extensors (rectus femoris [RF], vastus medialis [VM], vastus lateralis [VL]) using a Noraxon TeleMyo Desktop DTS System (Noraxon, Arizona, USA). The skin was prepared by shaving, abrading and cleansing with 70% alcohol. Bipolar Ag/Ag/Cl surface electrodes (2 cm inter-electrode distance, Noraxon) were attached over the belly of each muscle at SENIAM recommended recording sites (Stegeman & Hermans, 2007), parallel to the presumed orientation of the muscle fibres. The raw EMG signals were wirelessly transmitted (Wireless Research EMG Probes, Part 542, Noraxon) to a receiver (Desktop DTS, Part 586, Noraxon), amplified x500, sampled at 2000 Hz in synch with torque via the same A/D converter and PC software, and band-pass filtered off-line between 6 and 500 Hz using a fourth-order zero-lag Butterworth filter.

Muscle Architecture

A static ultrasound image (Hitachi Noblus, Hitachi Medical Systems, UK) of the VL was taken using a linear array probe with a 94 mm scan width (HI VISION L53L, Hitachi Medical Systems, UK). The image was taken prior to any other measurements whilst the participant

was seated in the dynamometer at rest, and with a joint angle of 100° (where 180° is full knee extension). The probe was placed perpendicular to the skin surface, over the thickest part of the belly of the VL, at 50% of the line between the greater trochanter and the knee joint centre, and aligned so that the muscle fascicles of the VL and their insertion into the deep aponeurosis were clearly visible.

Muscle thickness, pennation angle and fascicle length (Figure 1) were determined from the still images offline using Tracker software (an open source Video Analysis Tool, available from <http://physlets.org/tracker/>). Muscle thickness was defined as the mean distance between the deep and superficial aponeuroses at three points: at the middle and either end of the image. Pennation angle was defined as the mean of the angle between three separate muscle fascicles and their insertion on the deep aponeurosis. Fascicle length was extrapolated from the pennation angle and muscle thickness using trigonometry (de Brito Fontana, Roesler & Herzog 2014, Franchi et al. 2014, Tillin, Pain & Folland 2012), as the entire length of the fascicle was not visible in the image. Between-session reliability of muscle architecture measures was assessed in a pilot study of eight able-bodied controls using the same methods as described above. Coefficient of Variation (CV) was 4.4%, 10.9% and 9.3% for muscle thickness, fascicle length, and pennation angle, respectively.

Electrical Stimulation

Square wave (0.2 ms duration) electrical impulses were delivered percutaneously to the femoral nerve, via a constant current variable voltage stimulator (Model DS7AH, Digitimer, Ltd, Welwyn Garden City, UK), to evoke supramaximal twitch, doublet and octet contractions of the knee extensors. The cathode stimulation probe (1 cm diameter, protruding 2 cm from a plastic base, Electro Medical Supplies, Wantage UK) was firmly pressed into the femoral triangle in the position that evoked the greatest twitch response for a submaximal (30–60 mA) electrical current. The anode (10 x 7 cm carbon rubber electrode) was taped in place over the greater trochanter. Single impulses were delivered with step-wise increments in the

current, separated by 15 s, until a plateau in the amplitude of twitch torque and compound muscle action potentials (M-waves) were reached. The stimulus intensity was then increased by 20% to ensure supramaximal stimulation, and three supramaximal twitch contractions, separated by 20 s, were delivered. The current was reduced prior to commencing the octet contractions (eight pulses at 300 Hz), and stepwise increments in the current were delivered 15 s apart until the supramaximal current used for twitch contractions was attained. Subsequently, three supramaximal octet contractions were evoked.

The mean M-wave peak-to-peak amplitude of the three supramaximal twitch contractions was defined as the maximal M-wave (M_{\max}) for each muscle. Torque measurements from the evoked contractions were twitch and octet peak torque (PT) and peak RTD (calculated using a 15 ms moving time window) presented as absolute and relative to PT. These variables were averaged across the three supra-maximal twitch and octet contractions recorded.

Knee Extension MVCs

Participants performed a series of ~20 warm-up contractions of 3-s duration at progressively higher intensities before completing six maximum voluntary contractions (MVCs). Each MVC lasted 3-5 s and was followed by 30-60 s rest. Participants were instructed to push 'as hard as possible' and strong verbal encouragement was given throughout the contractions. Real-time biofeedback of torque output was provided on a computer monitor in front of participants. MVT was defined as the greatest instantaneous peak voluntary torque (not due to superimposed stimulation) recorded during any of the MVCs or explosive contractions.

Explosive Voluntary Contractions

Participants completed 10-15 explosive isometric contractions, each separated by 20 s rest, utilising the method described by Folland, Buckthorpe & Hannah (2014). Three explosive voluntary contractions were chosen for analysis and all dependent variables assessed were averaged across these three explosive contractions. The three contractions were chosen as

those with the highest peak RTD, peak torque >80% MVT, and no visible countermovement or pre-tension (quantified as change of baseline torque <0.5 Nm during the 100 ms prior to visible torque onset) were used for analysis. Peak RTD was extracted and expressed as both an absolute and relative to MVT.

To assess neural drive during the explosive contractions, the RMS amplitude of the EMG signal for each quadriceps muscle was calculated for the time period 0-100 ms from EMG onset (EMG_{0-100}), and normalised to M_{max} at the same muscle before averaging across the three quadriceps muscles. EMG onsets, defined as the onset of the first muscle to be activated, were identified with a standardised systematic protocol of visual identification (Tillin et al. 2010).

Voluntary Activation

The 2nd, 4th, and 6th MVCs had a single doublet superimposed at the plateau of the torque-time curve, and two further doublets evoked at rest immediately after the MVC. The difference between superimposed and resting potentiated doublet torque was used to VA (a measure of neural drive at MVT), using the equation:

$$VA(\%) = 100 \times (1 - D_s/D_c)$$

(1)

where D_s and D_c are the superimposed and control doublets, respectively.

The root mean square (RMS) of the EMG signal for each quadriceps muscle was calculated over the 500 ms window centred on or nearest to MVT, which was not influenced by stimulation artefact (EMG_{MVT}). EMG_{MVT} was normalised to M_{max} of the same muscle and averaged across the three quadriceps muscles.

Walking Gait

Kinematic data were collected using twelve Vicon Vantage V5 (Vicon Motion Systems Ltd.; Oxford, UK) motion capture cameras sampling at 200 Hz synched with three in-series Kistler force plates (Type 9281c; Kistler Instruments Ltd., Hampshire, UK) in the middle of a 15 m walkway sampling force data at 1000 Hz. Two sets of Brower TC timing gates (Brower Timing, Utah, USA) placed 2 m either side of the force plates were used to capture average walking pace. Retroreflective markers (14 mm diameter) were placed on the skin according to the Plug-In-Gait lower-body marker set. Markers for the shank, ankle and foot were placed in positions on the prosthetic corresponding as closely as possible to those on the intact limb.

Data collection involved participants walking along the 15 m walkway at a self-selected, habitual pace. Average walking pace was determined in preliminary trials by allowing participants to walk up and down the walkway until speed stabilised. Three 'good' trials, defined as a single pass with a successful force plate strike, walking speed within $\pm 5\%$ of average, and no gaps in marker data greater than 40 frames, were selected for analysis for each limb. Data were processed in Vicon Nexus 2.7.1. Raw marker trajectories and analogue force data were filtered using a low-pass zero-lag fourth-order Butterworth filter, at cut-off frequencies of 8 and 200 Hz respectively. Standard inverse dynamics techniques were used to calculate net internal joint moments, normalised by body mass (Winter and Sienko 1988).

Internal peak knee extension moment and total impulse (calculated as the integral of internal knee extension moment with respect to time) for the entire stance phase were extracted for each limb and averaged across the trials selected for analysis.

Statistical Analysis

Paired t-tests revealed no differences in either MVT or peak RTD between dominant vs. non-dominant (MVT: $p = 0.775$, $g = 0.07$; RTD: $p = 0.237$, $g = 0.43$) limbs in the control group, where the dominant limb was defined as the one in which the participant would favour

to kick a ball. Thus, each dependent variable was averaged between the dominant and non-dominant limbs in the control group, and comparisons are made between the mean of the control limbs (CON) vs the amputated limb of ITTAs (AMP) vs the intact limb (INT).

Levene's test was used to check for equality of variances prior to running all analyses. A one-way mixed design ANOVA was used to analyse the influence of limb (AMP vs. INT vs. CON) on each dependent variable. In the instance of a main effect for any of the ANOVAs, post-hoc Bonferroni corrected t-tests (paired t-tests for AMP vs. INT, and independent t-tests for AMP or INT vs. CON) were used for paired comparisons. Effect size, Hedges g , was calculated for paired comparisons, and interpreted as small (0.2-0.5), medium (0.5-0.8) and large effects (>0.8). Statistical analysis was completed using SPSS version 24, and the significance level was set at $p < 0.05$. Data are reported as mean \pm standard deviation (SD), with absolute percentage difference in values between each condition.

Results

Due to an injury to one ITTA occurring between visits 2 and 3 (neuromuscular and gait assessment) data are for 9 and 8 ITTAs, respectively. One control withdrew from octet and doublet stimulation, so control data for VA and octet variables are presented for 8 controls, but all other variables are for 9 controls. The groups had similar age, height, body mass and physical activity scores ($p \geq 0.354$; $g = 0.10$ - 0.64 ; Table 1). There was a large effect size for the controls to walk faster ($g = 1.21$), although this difference was not statistically significant ($p = 0.616$; Table 1).

Muscle Architecture

There was no main effect ($p = 0.226$) of limb on pennation angle (Table 2). However, muscle thickness in AMP was lower than both INT (-41% , $p = 0.030$, $g = 1.78$) and CON (-38% , $p = 0.002$, $g = 1.58$; Figure 1), but similar between INT and CON ($p = 1.000$, $g = 0.23$; Table 2).

305 Fascicle length was shorter in AMP than INT (-36%, $p < 0.001$, $g = 0.95$), but similar
306 between AMP and CON ($p = 0.187$; $g = 0.50$), and INT and CON ($p = 1.000$; $g = 0.49$).

307 Contractile Properties

308 PT and absolute RTD in both the twitch and octet (Table 2) were lower in AMP than both
309 INT and CON (-72% to -50%, $p = 0.001$ -0.004, $g = 1.97$ -2.84), but similar between INT and
310 CON ($p \geq 0.284$, $g = 0.40$ -0.68).

311 When expressed relative to PT, twitch RTD was 18% lower ($p = 0.006$, $g = 1.35$), and octet
312 RTD 25% lower ($p < 0.001$, $g = 2.60$) in AMP when compared to INT (Table 2). Relative
313 twitch and octet RTD were also both 14% lower in AMP compared to CON (twitch RTD: $p =$
314 0.036, $g = 1.59$; octet RTD: $p = 0.037$, $g = 1.63$). Despite being statistically similar, there was
315 a large effect for relative octet RTD to be greater in INT than CON ($p = 0.120$, $g = 1.03$;
316 Table 2), whilst relative twitch RTD was similar between INT and CON ($p = 1.000$, $g = 0.18$).

317 Maximal and Voluntary Explosive Torque

318 MVT (both absolute and relative to body mass) was significantly lower in AMP compared to
319 both INT (~-60%, $p < 0.002$, $g = 1.74$ -1.97) and CON (~-64%, $p < 0.001$, $g = 2.05$ -2.33).
320 There were no differences between INT and CON in absolute ($p = 1.000$, $g = 0.35$) or
321 relative MVT ($p = 1.000$, $g = 0.28$; Figure 2A and C).

322 Absolute peak voluntary RTD (Figure 2B) was ~75% lower in AMP than INT ($p = 0.001$, $g =$
323 2.22), ~76% lower in AMP than CON ($p < 0.001$, $g = 2.36$), but similar between INT and
324 CON ($p = 1.000$, $g = 0.14$). Relative to MVT, peak RTD was significantly smaller in AMP
325 than INT (-43%, $p = 0.027$, $g = 1.37$) and CON (-39%, $p = 0.031$, $g = 1.09$), while INT and
326 CON were similar ($p = 1.000$, $g = 0.23$; Figure 2D).

327 Neural Drive

Both VA and RMS EMG_{MVT} (Table 2) were lower in AMP than INT (-44% for VA, $p < 0.001$, $g = 3.63$; and -43% for EMG_{MVT}, $p < 0.001$, $g = 1.97$) and CON (-43% for VA, $p < 0.001$, $g = 3.54$; -32% for EMG_{MVT}, $p = 0.021$, $g = 1.23$), but similar between INT and CON ($p \geq 0.271$, $g = 0.14-0.70$).

During the voluntary explosive contractions, there was no main effect of limb ($p = 0.304$) on the amplitude of explosive RMS EMG₀₋₁₀₀ (Table 2). However, there was a moderate effect for RMS EMG₀₋₁₀₀ to be greater in INT than AMP ($g = 0.75$), but only small to moderate effects for other comparisons (AMP vs. CON, $g = 0.30$; INT vs. CON, $g = 0.45$).

Knee Kinetics in Gait

Knee moments throughout stance are presented in Figure 3. Both absolute and relative peak knee extensor moment during the stance phase of gait was significantly lower in the AMP compared to INT (absolute -59%, $p = 0.011$, $g = 1.77$; BM -60%, $p = 0.005$, $g = 1.78$) and CON (absolute -54%, $p = 0.005$, $g = 1.61$; BM -59%, $p = 0.006$, $g = 1.72$) limbs, but similar between INT and CON ($p = 1.000$; $g = 0.05-0.14$; Table 2). While there was no main effect of limb on absolute or relative knee extensor moment impulse during stance ($p > 0.069$), there were medium to large effects for it to be lower in AMP than INT (-36%; absolute $g = 0.99$, BM $g = 1.15$) and CON (-27%; absolute $g = 0.56$, BM $g = 0.90$), respectively (Table 2).

Discussion

In this study, we compared quadriceps strength and neuromuscular function in the amputated limb of ITTAs with their intact limb and a control group limb, providing a novel model for studying the long-term (>1.5 years) effects of chronic disuse. Long-term disuse of the amputated limb in ITTAs was evidenced from the ~60% lower peak knee extensor moments during walking compared to the intact and control limbs. This disuse was accompanied by ~60% lower MVT and ~75% lower RTD in the amputated limb, which are

much greater differences than may be predicted from short-term disuse studies. Declines in MVT appeared to be largely due to reduced muscle size (evidenced by lower muscle thickness in AMP) and neural drive (evidenced by lower VA and EMG_{MVT} in AMP). Declines in RTD appeared to be due primarily to declines in MVT and a shift towards slower intrinsic contractile properties, with neural drive in explosive contractions being unaffected in AMP.

TTAs as a model for long-term disuse

In the current study, there were large effects for knee extensor kinetics during gait to be lower in amputated than intact or control limbs which, coupled with the considerable reductions in knee extensor strength in the amputated limb, suggests the amputated limb undergoes substantially less habitual loading during ambulation. The reduced knee extensor moments in gait may also be partly due to increased co-contraction at the knee on the amputated limb during gait (Culham et al., 1986; Isakov et al., 2001). Future research should therefore aim to quantify internal loading of the knee extensors for a more direct estimation of disuse and its association with strength changes in the amputated limb. Consistent with our results, previous studies have reported decreased knee moments (Powers, Rao & Perry, 1998, Winter & Sienko, 1988); powers (Powers, Rao & Perry, 1998, Winter & Sienko, 1988); and work (Silverman & Neptune, 2012) on the amputated limb in walking. In contrast to these previous studies however, the ITTAs of the current study were young, healthy, and moderate-highly active. As a result, the effects of the evident disuse on strength and neuromuscular function could be isolated from factors such as ageing, disease, and sedentary behaviour, which are known to independently affect muscle strength and function (Narici & de Boer 2011, Sacchetti et al. 2013).

The knee extensors of the intact limb in the ITTAs did not differ from those of an able-bodied control population for kinetics during walking, MVT, RTD, or any of the neuromuscular determinants of strength. This suggests that, for these parameters, the intact limb of the

ITTAs provides an ideal internal control for comparison to the amputated limb, from which to draw conclusions about the effects of chronic disuse.

Changes in Strength

The declines in MVT found in the amputated limb compared to the intact limb (-59%) are comparable to, albeit at the high end of, differences observed in previous amputee studies (-33 to -57%; Isakov et al. 1996, Lloyd et al. 2010, Moirenfeld et al. 2000, Pedrinelli et al. 2002), but considerably greater than the reduction in strength typically observed after a period of short-term disuse of up to 120 days (~23%; Narici & de Boer, 2011). Short-term intervention studies suggest that MVT decreases exponentially over time following unloading, plateauing out after ~90 days; however, the results of this study suggest the strength declines with longer-term disuse are considerably more than could be predicted from short-term intervention studies.

To the authors' knowledge, only two previous studies have investigated the effect of disuse on voluntary RTD of the knee extensors, reporting 54% (Bamman et al. 1998) and 42% (de Boer et al. 2007) decreases in RTD after 16 days of bed rest, and 23 days of ULLS, respectively. The considerable reductions in peak RTD (-75%) in the amputated vs. intact limb are important, as RTD is considered more functionally relevant than MVT, in many sports-specific and daily tasks, such as sprinting, jumping, and balance recovery (Behan, Pain & Folland 2018, Pijnappels et al. 2008, Tillin, Pain & Folland 2013).

When expressed relative to MVT, peak RTD was significantly reduced in the amputated compared to the non-amputated limbs, although limb differences were considerably smaller for relative than absolute peak RTD. Thus, the reduction in MVT appears to be a large contributing factor to reduced absolute RTD in the amputated limb; however, this only partially contributed to the reduction in peak RTD, which was likely also influenced by the slowing of the contractile properties (discussed in more detail below).

Mechanisms of Strength Differences

Neural Drive

A broad suppression in neuromuscular activity at maximal force production – indicated by reduced amputated limb VA (~44%) and EMG_{MVT} (~38%) compared to non-amputated limbs – likely contributes to the reduction in amputated limb MVT. Whilst previous studies have reported reduced quadriceps EMG amplitude (-16 to -35%; Alkner & Tesch 2004, Deschenes et al. 2002) and VA (-7%; Kawakami et al. 2001), others have not observed changes in these measurements (de Boer et al. 2007, Campbell et al. 2013, Horstman et al. 2012), following periods of disuse of up to 89 days. Thus, the large limb effects on VA and EMG responses observed in the present study suggest that reductions in neural drive with disuse become more pronounced and observable over time. Of note is the specificity of the neural deficits in the ITTAs to the amputated limb. Evidence from unilateral injury and training studies suggest a cross-over effect of neural function, in that neural drive adaptations occur at the contralateral, as well as the injured/trained limb (Bogdanis et al. 2019, Tillin et al. 2011). In this study however, there was no evidence that reduced neural drive on the amputated side had affected neural drive on the intact side, which was similar to the control limb. Perhaps this is because ITTAs rely more on the intact limb for most activities of daily living and exercise (e.g. Winter & Sienko 1988), which may negate any cross-over effects of reduced neural drive from the amputated to the intact limb.

Despite the substantial differences between the amputated and non-amputated limbs evident in neural drive during maximum force production, no such differences were observed in this study in explosive-phase EMG amplitude (Table 2). This suggests that altered neural drive does not explain the lower peak RTD in the amputated limb, which is interesting given that neural drive is a key determinant of RTD (del Vecchio et al. 2019, Folland, Buckthorpe & Hannah 2014). The large variability in EMG, even after normalisation to M_{max} (Buckthorpe et al. 2012), greater variability in RTD compared to MVT (Folland, Buckthorpe & Hannah 2014,

Tillin, Pain & Folland 2013), and small sample sizes ($n = 9$ per limb) may have reduced the chances of observing a significant effect. Alternatively, the amputated limb's role in ambulation may explain the lack of differences in neural drive during the explosive contractions. Specifically, whilst the knee extensors of the amputated limb experience reduced load compared to the intact during ambulation, the amputated side does contribute to stability and postural correction, for which RTD appears to be important (Behan, Pain & Folland 2018). Thus, typical physical activity in the amputees may provide sufficient stimulus to maintain the neural drive during short, rapid contractions, which typically underpins RTD.

Muscle Architecture

The VL muscle was 41% thinner in the amputated limb compared to the intact, which is a larger difference than the declines in MRI and CT scanner measurements of muscle size (-3 to -18%) observed in short-term disuse studies (Alkner & Tesch 2004, Campbell et al. 2013, de Boer et al. 2007, Dirks et al. 2016). Thus, similar to the changes observed for strength and neural drive, reductions in muscle size with long-term disuse are much greater than could be predicted from short-term disuse studies. Muscle size is considered an important determinant of MVT (Blazevich et al. 2009), and thus the reduction in muscle thickness is likely to contribute to the declines in both MVT, and by association RTD, in the amputated limb.

Fascicle length was reduced by 36% in the amputated limb compared to the intact. Again, this difference is considerably greater than the decline in knee extensor fascicle length (6-9%) typically observed with short-term unloading (Campbell et al. 2013, de Boer et al. 2007). ITTAs walk with a comparatively stiff knee joint on the amputated limb (Powers, Rao & Perry 1998, Winter & Sienko 1988), which would theoretically isolate loading to shorter fascicle lengths, and limit the stimulus likely required to maintain longer fascicle lengths. Decreases in fascicle length may reduce maximum shortening velocities and power (Blazevich & Sharp 2005), and shift the torque-angle relationship towards more extended knee positions

(Blazevich et al. 2009). Given our strength measurements were made at a typical plateau region of the torque-angle relationship (Chow et al. 1999), a shift away from this region in the amputated limb may have partly contributed to the observed differences in MVT and RTD.

In contrast to the results of previous research which demonstrated decreases in pennation angle during short periods of ULLS (de Boer et al. 2007, Campbell et al. 2013), our results appear to suggest that pennation angle does not change with long-term disuse. In healthy populations, angles of pennation of the VL muscle have been reported to be 6-27° (Blazevich et al. 2006, Rutherford & Jones 1992); the pennation angle of all three limbs in this study (~12-14°) falls within this range. This suggests that the structural re-modelling that seems to take place in the early phases of disuse are not representative of long-term adaptations. It is possible muscle thickness declines at a faster rate than fascicle length with short term disuse, causing a decline in pennation angle; whilst over longer periods of disuse, fascicle length reductions “catch-up” with muscle thickness loss, causing a return to baseline pennation angle, but this hypothesis cannot be tested with our data.

Intrinsic contractile properties

The significant reductions in evoked (twitch and octet) contractile peak torque in the amputated compared to the intact and control limbs (Table 2) are reflective of the reduced capacity of the amputated limb knee extensors for torque production. These changes were accompanied by reductions in RTD, both absolute and relative to peak torque, reflecting a shift towards slower contractile properties in the intact limb. This is in contrast to the results of short-term disuse studies in both healthy controls and pathological populations, which have reported a shift towards faster contractile properties owing to a greater expression of fast-contracting myosin-heavy-chain isoforms (MHC; Bamman et al. 1998, Kapchisky et al. 2018, Trappe et al. 2004). The results of the current study therefore provide novel evidence that changes in intrinsic contractile properties with long-term disuse are more characteristic

of ageing muscle, which also displays a slowing of the contractile properties (Roos et al. 1999). This slowing may be due to preferential atrophy of type II muscle fibres, and potentially also to an increased dominance of type I MHC in fibres co-expressing MHCs commonly seen with old age (Lexell et al. 1988). The slower contractile properties in the amputated limb likely contributed to the lower voluntary peak RTD also observed in the amputated limb, as twitch and octet RTD are important determinants of voluntary RTD (Folland, Buckthorpe & Hannah 2014).

Conclusion

This study was the first to utilise ITTAs as a novel study model to investigate the effects of long-term muscle disuse on strength and neuromuscular function, in young, healthy, active adults. Strength, neuromuscular function and loading during gait, of the intact limb of ITTAs were comparable to a control able-bodied limb, suggesting the intact limb provides a suitable internal control for comparison to the amputated limb for these parameters. The quadriceps muscles of the amputated limb displayed considerably less habitual loading during gait, than the intact side. This disuse of the amputated limb was accompanied by larger reductions in MVT and RTD than could be predicted from short-term disuse studies. The reductions in MVT were likely due to the declines in muscle size and neural drive, whilst the reductions in RTD appeared due to the decline in MVT coupled with a slowing of the contractile properties.

501

References

- 502 Alkner, B. A., & Tesch, P. A. (2004). Knee extensor and plantar flexor muscle size and
503 function following 90 days of bed rest with or without resistance exercise. *European*
504 *Journal of Applied Physiology*, 93(3), 294-305. doi: 10.1007/s00421-004-1172-8
- 505 Balshaw, T.G., Massey, G.J., Maden-Wilkinson, T.M., Tillin, N.A. and Folland, J.P., 2016.
506 Training-specific functional, neural, and hypertrophic adaptations to explosive-vs.
507 sustained-contraction strength training. *Journal of Applied Physiology*, 120(11),
508 pp.1364-1373. doi: 10.1152/jappphysiol.00091.2016
- 509 Bamman, M. M., Clarke, M. S., Feedback, D. L., Talmadge, R. J., Stevens, B. R., Lieberman,
510 S. A., & Greenisen, M. C. (1998). Impact of resistance exercise during bed rest on
511 skeletal muscle sarcopenia and myosin isoform distribution. *Journal of Applied*
512 *Physiology*, 84(1), 157-163. doi: 10.1152/jappl.1998.84.1.157
- 513 Bassey, E. J., Fiatarone, M. A., O'Neill, E. F., Kelly, M., Evans, W. J., & Lipsitz, L. A. (1992).
514 Leg extensor power and functional performance in very old men and women. *Clinical*
515 *Science*, 82(3), 321-327. doi: 10.1042/cs0820321
- 516 Behan, F. P., Pain, M. T., & Folland, J. P. (2018). Explosive voluntary torque is related to
517 whole-body response to unexpected perturbations. *Journal of Biomechanics*, 81, 86-
518 92. doi: 10.1016/j.jbiomech.2018.09.016
- 519 Berg, H. E., & Tesch, P. A. (1996). Changes in muscle function in response to 10 days of
520 lower limb unloading in humans. *Acta Physiologica*, 157(1), 63-70. doi: 10.1046/j.1365-
521 201x.1996.476217000.x
- 522 Blazeovich, A. J., Cannavan, D., Horne, S., Coleman, D. R., & Aagaard, P. (2009). Changes
523 in muscle force-length properties affect the early rise of force in vivo. *Muscle & Nerve*,
524 39(4), 512-520. doi: 10.1002/mus.21259

525 Blazeovich, A. J., Gill, N. D., & Zhou, S. (2006). Intra-and intermuscular variation in human
 526 quadriceps femoris architecture assessed in vivo. *Journal of Anatomy*, 209(3), 289-
 527 310. doi: 10.1111/j.1469-7580.2006.00619.x

528 Blazeovich, A. J., & Sharp, N. C. (2005). Understanding muscle architectural adaptation:
 529 macro-and micro-level research. *Cells Tissues Organs*, 181(1), 1-10. doi:
 530 10.1159/000089964

531 Bogdanis, G. C., Tsoukos, A., Kaloheri, O., Terzis, G., Veligekas, P., & Brown, L. E. (2019).
 532 Comparison between unilateral and bilateral plyometric training on single-and double-
 533 leg jumping performance and strength. *The Journal of Strength & Conditioning*
 534 *Research*, 33(3), 633-640. doi: 10.1519/jsc.0000000000001962

535 Brown, C. J., Friedkin, R. J., & Inouye, S. K. (2004). Prevalence and outcomes of low
 536 mobility in hospitalized older patients. *Journal of the American Geriatrics Society*,
 537 52(8), 1263-1270. doi: 10.1111/j.1532-5415.2004.52354.x

538 Buckthorpe, M. W., Hannah, R., Pain, T. G., & Folland, J. P. (2012). Reliability of
 539 neuromuscular measurements during explosive isometric contractions, with special
 540 reference to electromyography normalization techniques. *Muscle & Nerve*, 46(4), 566-
 541 576. doi:10.1002/mus.23322

542 Campbell, E., Seynnes, O. R., Bottinelli, R., McPhee, J. S., Atherton, P. J., Jones, D. A.,
 543 Narici, M. V. (2013). Skeletal muscle adaptations to physical inactivity and subsequent
 544 retraining in young men. *Biogerontology*, 14(3), 247-259. doi: 10.1007/s10522-013-
 545 9427-6

546 Chow, J. W., Darling, W. G., & Ehrhardt, J. C. (1999). Determining the force-length-velocity
 547 relations of the quadriceps muscles: II. Maximum muscle stress. *Journal of Applied*
 548 *Biomechanics*, 15(2), 191-199. doi: 10.1123/jab.15.2.191

549 Dirks, M.L., Wall, B.T., van de Valk, B., Holloway, T.M., Holloway, G.P., Chabowski, A.,
 550 Goossens, G.H. and van Loon, L.J., 2016. One week of bed rest leads to substantial

551 muscle atrophy and induces whole-body insulin resistance in the absence of skeletal
552 muscle lipid accumulation. *Diabetes*, 65(10), 2862-2875. doi: 10.2337/db15-1661

553 de Boer, M. D., Maganaris, C. N., Seynnes, O. R., Rennie, M. J., & Narici, M. V. (2007).
554 Time course of muscular, neural and tendinous adaptations to 23 day unilateral lower-
555 limb suspension in young men. *The Journal of Physiology*, 583(3), 1079-1091. doi:
556 10.1113/jphysiol.2007.135392

557 de Brito Fontana, H., Roesler, H. and Herzog, W., 2014. In vivo vastus lateralis force–
558 velocity relationship at the fascicle and muscle tendon unit level. *Journal of*
559 *Electromyography and Kinesiology*, 24(6), pp.934-940. doi:
560 10.1016/j.jelekin.2014.06.010

561 Deschenes, M. R., Giles, J. A., McCoy, R. W., Volek, J. S., Gomez, A. L., & Kraemer, W. J.
562 (2002). Neural factors account for strength decrements observed after short-term
563 muscle unloading. *American Journal of Physiology-Regulatory, Integrative and*
564 *Comparative Physiology*, 282(2), R583. doi: 10.1152/ajpregu.00386.2001

565 Fey, N. P., Silverman, A. K., & Neptune, R. R. (2010). The influence of increasing steady-
566 state walking speed on muscle activity in below-knee amputees. *Journal of*
567 *electromyography and Kinesiology*, 20(1), 155-161. doi: 10.1016/j.jelekin.2009.02.004

568 Folland, J. P., Buckthorpe, M. W., & Hannah, R. (2014). Human capacity for explosive force
569 production: Neural and contractile determinants. *Scandinavian Journal of Medicine &*
570 *Science in Sports*, 24(6), 894-906. doi: 10.1111/sms.12131

571 Franchi, M.V., Atherton, P.J., Reeves, N.D., Flück, M., Williams, J., Mitchell, W.K., Selby, A.,
572 Beltran Valls, R.M. & Narici, M.V., (2014). Architectural, functional and molecular
573 responses to concentric and eccentric loading in human skeletal muscle. *Acta*
574 *Physiologica*, 210(3), pp.642-654. doi: 10.1111/apha.12225

575 Horstman, A. M., Ruiter, C. J. d., Duijnhoven, N T L van, Hopman, M. T. E., & Haan, A. D.
576 (2012). Changes in muscle contractile characteristics and jump height following 24

577 days of unilateral lower limb suspension. *European Journal of Applied Physiology*,
578 112(1), 135-144. doi:10.1007/s00421-011-1958-4

579 Isakov, E., Burger, H., Gregorič, M., & Marinček, C. (1996). Isokinetic and isometric strength
580 of the thigh muscles in below-knee amputees. *Clinical Biomechanics*, 11(4), 233-235.
581 doi: 10.1016/0268-0033(95)00078-x

582 Kawakami, Y., Akima, H., Kubo, K., Muraoka, Y., Hasegawa, H., Kouzaki, M., & Fukunaga,
583 T. (2001). Changes in muscle size, architecture, and neural activation after 20 days of
584 bed rest with and without resistance exercise. *European journal of applied physiology*,
585 84(1-2), 7-12. doi: 10.1007/s004210000330

586 Kapchinsky, S., Vuda, M., Miguez, K., Elrief, D., de Souza, A.R., Bagloli, C.J., Aare, S.,
587 MacMillan, N.J., Baril, J., Rozakis, P. and Sonjak, V., 2018. Smoke-induced
588 neuromuscular junction degeneration precedes the fibre type shift and atrophy in
589 chronic obstructive pulmonary disease. *The Journal of physiology*, 596(14), pp.2865-
590 2881.

591 Lambertz, D., Pérot, C., Kaspranski, R., & Goubel, F. (2001). Effects of long-term spaceflight
592 on mechanical properties of muscles in humans. *Journal of Applied Physiology*, 90(1),
593 179-188. doi: 10.1152/jappl.2001.90.1.179

594 Lexell, J., Taylor, C. C., & Sjöström, M. (1988). What is the cause of the ageing atrophy?:
595 Total number, size and proportion of different fiber types studied in whole vastus
596 lateralis muscle from 15-to 83-year-old men. *Journal of the Neurological Sciences*,
597 84(2-3), 275-294.

598 Lloyd, C. H., Stanhope, S. J., Davis, I. S., & Royer, T. D. (2010). Strength asymmetry and
599 osteoarthritis risk factors in unilateral trans-tibial, amputee gait. *Gait & Posture*, 32(3),
600 296-300. doi: 10.1016/j.gaitpost.2010.05.003

601 Moirenfeld, I., Ayalon, M., Ben-Sira, D., & Isakov, E. (2000). Isokinetic strength and
602 endurance of the knee extensors and flexors in trans-tibial amputees. *Prosthetics and*
603 *Orthotics International*, 24(3), 221-225. doi: 10.1080/03093640008726551

604 Narici, M. V., & De Boer, M. D. (2011). Disuse of the musculo-skeletal system in space and
605 on earth. *European Journal of Applied Physiology*, 111(3), 403-420. doi:
606 10.1007/s00421-010-1556-x

607 Nolan, L., Wit, A., Dudziński, K., Lees, A., Lake, M. and Wychowański, M., 2003.
608 Adjustments in gait symmetry with walking speed in trans-femoral and trans-tibial
609 amputees. *Gait & Posture*, 17(2), pp.142-151. doi: 10.1016/S0966-6362(02)00066-8

610 Pedrinelli, A., Saito, M., Coelho, R. F., Fontes, R., & Guarniero, R. (2002). Comparative
611 study of the strength of the flexor and extensor muscles of the knee through isokinetic
612 evaluation in normal subjects and patients subjected to trans-tibial amputation.
613 *Prosthetics and Orthotics International*, 26(3), 195-205. doi:
614 10.1080/03093640208726648

615 Pijnappels, M., Reeves, N. D., Maganaris, C. N., & Van Dieen, J. H. (2008). Tripping without
616 falling; lower limb strength, a limitation for balance recovery and a target for training in
617 the elderly. *Journal of Electromyography and Kinesiology*, 18(2), 188-196. doi:
618 10.1016/j.jelekin.2007.06.004

619 Powers, C. M., Boyd, L. A., Fontaine, C. A., & Perry, J. (1996). The influence of lower-
620 extremity muscle force on gait characteristics in individuals with below-knee
621 amputations secondary to vascular disease. *Physical Therapy*, 76(4), 85. doi:
622 10.1093/ptj/76.4.369

623 Powers, C. M., Rao, S., & Perry, J. (1998). Knee kinetics in trans-tibial amputee gait. *Gait &*
624 *Posture*, 8(1), 1-7. doi: 10.1016/s0966-6362(98)90210-7

625 Puthucheary, Z. A., McNelly, A. S., Rawal, J., Connolly, B., Sidhu, P. S., Rowlerson, A., ...
626 Montgomery, H. E. (2017). Rectus Femoris Cross-Sectional Area and Muscle Layer

627 Thickness: Comparative Markers of Muscle Wasting and Weakness. *American Journal*
628 *of Respiratory and Critical Care Medicine*, 195(1), 136–138. doi:10.1164/rccm.201604-
629 0875LE

630 Renström, P., Grimby, G. and Larsson, E., 1983. Thigh muscle strength in below-knee
631 amputees. *Scandinavian Journal of Rehabilitation Medicine. Supplement*, 9, 163-173.

632 Roos, M. R., Rice, C. L., Connelly, D. M., & Vandervoort, A. A. (1999). Quadriceps muscle
633 strength, contractile properties, and motor unit firing rates in young and old men.
634 *Muscle & Nerve*, 22(8), 1094-1103. doi: 10.1002/(sici)1097-
635 4598(199908)22:8<1094::aid-mus14>3.0.co;2-g

636 Rozier, C. K., Elder, J. D., & Brown, M. (1979). Prevention of atrophy by isometric exercise
637 of a casted leg. *The Journal of sports medicine and physical fitness*, 19(2), 191.

638 Rutherford, O. M., & Jones, D. A. (1992). Measurement of fibre pennation using ultrasound
639 in the human quadriceps in vivo. *European Journal of Applied Physiology and*
640 *Occupational Physiology*, 65(5), 433-437. doi: 10.1007/bf00243510

641 Sacchetti, M., Balducci, S., Bazzucchi, I., Carlucci, F., Scotto di Palumbo, A., Haxhi, J.,
642 Pugliese, G. (2013). Neuromuscular dysfunction in diabetes: Role of nerve impairment
643 and training status. *Med Sci Sports Exerc*, 45(1), 52-59. doi:
644 10.1249/mss.0b013e318269f9bb

645 Sanderson, D.J. & Martin, P.E., 1997. Lower extremity kinematic and kinetic adaptations in
646 unilateral below-knee amputees during walking. *Gait & posture*, 6(2), pp.126-136. doi:
647 10.1016/S0966-6362(97)01112-0

648 Schmalz, T., Blumentritt, S., & Marx, B. (2007). Biomechanical analysis of stair ambulation in
649 lower limb amputees. *Gait & Posture*, 25(2), 267-278. doi:
650 10.1016/j.gaitpost.2006.04.008

651 Schoeman, M., Diss, C. E., & Strike, S. C. (2012). Kinetic and kinematic compensations in
652 amputee vertical jumping. *Journal of Applied Biomechanics*, 28(4), 438-447. doi:
653 10.1123/jab.28.4.438

654 Silverman, A. K., & Neptune, R. R. (2012). Muscle and prosthesis contributions to amputee
655 walking mechanics: A modeling study. *Journal of Biomechanics*, 45(13), 2271-2278.
656 doi: 10.1016/j.jbiomech.2012.06.008

657 Stegeman, D. F., & Hermens, H. J. (2007). Standards for surface electromyography: The
658 european project surface EMG for non-invasive assessment of muscles (SENIAM).
659 *Enschede: Roessingh Research and Development*, 108-112.

660 Tillin, N. A., Jimenez-Reyes, P., Pain, M. T., & Folland, J. P. (2010). Neuromuscular
661 performance of explosive power athletes versus untrained individuals. *Medicine and*
662 *Science in Sports and Exercise*, 42 (4), 781-790. doi:
663 10.1249/MSS.0b013e3181be9c7e

664 Tillin, N. A., Pain, M. T., & Folland, J. P. (2011). Short-term unilateral resistance training
665 affects the agonist–antagonist but not the force–agonist activation relationship. *Muscle*
666 *& Nerve*, 43(3), 375-384. doi: 10.1002/mus.21885

667 Tillin, N. A., Pain, M. T. & Folland, J. P. (2012). Short-term training for explosive strength
668 causes neural and mechanical adaptations. *Experimental Physiology*, 97: 630-641.
669 doi:10.1113/expphysiol.2011.063040

670 Tillin, N. A., Pain, M. T. G., & Folland, J. (2013). Explosive force production during isometric
671 squats correlates with athletic performance in rugby union players. *Journal of Sports*
672 *Sciences*, 31(1), 66-76. doi: 0.1080/02640414.2012.720704

673 Trappe, S., Trappe, T., Gallagher, P., Harber, M., Alkner, B., & Tesch, P. (2004). Human
674 single muscle fibre function with 84 day bed-rest and resistance exercise. *The Journal*
675 *of Physiology*, 557(2), 501-513. doi: 10.1113/jphysiol.2004.062166

676 Vecchio, A. D., Negro, F., Holobar, A., Casolo, A., Folland, J. P., Felici, F., & Farina, D.
677 (2019). You are as fast as your motor neurons: Speed of recruitment and maximal
678 discharge of motor neurons determine the maximal rate of force development in
679 humans. *The Journal of Physiology*, 597(9), 2445-2456. doi: 10.1113/JP277396

680 Winter, D. A., & Sienko, S. E. (1988). Biomechanics of below-knee amputee gait. *Journal of*
681 *Biomechanics*, 21(5), 361-367. doi: 10.1016/0021-9290(88)90142-x

682

683 **Author Contributions:** A.S., N.T., S.S. and S.M. conceived and designed the
684 research; A.S and S.M. performed experiments; A.S. analysed data; A.S. and N.T.
685 interpreted results of experiments; A.S. and N.T. drafted manuscript; A.S., N.T., S.S.
686 and S.M. edited and revised manuscript; all authors approved final version of the
687 manuscript. Data collection was performed in the biomechanics laboratory at the
688 University of Roehampton, Whitelands College.

689 **Acknowledgements:** The authors thank Alison Carlisle for her assistance during
690 testing and participants for their time in taking part in the study.

691 **Conflict of Interest Disclosure:** None.

692 **Funding:** This study was supported financially by the University of Roehampton's
693 Vice-Chancellor's Doctoral Scholarship.

694

695
696

Tables

Table 1. Participant information. Data are mean ± SD, and presented for n = 9 for both groups, except walking speed (n = 8 for ITTAs and n = 9 for controls). Cause of amputation was trauma for all ITTAs.

	ITTAs		Controls	
	Mean ± SD	Range	Mean ± SD	Range
Age (years)	40.3 ± 8.5	24 – 48	38.6 ± 6.3	27 – 46
Height (cm)	179 ± 8.2	165 – 186	177 ± 4.1	171 – 184
Body Mass (kg)	84.7 ± 16.7	54.6 – 114	80.0 ± 10.5	58.3 – 97.5
Activity Level (MET·min·week ⁻¹)	7890 ± 6122	480 – 15918	5686 ± 3256	2577 – 11817
Walking Speed (m·s ⁻¹)	1.34 ± 0.16	1.05 – 1.61	1.51 ± 0.10	1.34 – 1.71
Years since Amputation	12.2 ± 11.5	1.5 – 29.0	-	-

697

Table 2. Knee extensor kinetics and neuromuscular determinants of strength in the amputated (AMP) and intact (INT) limbs of unilateral transtibial amputees, and in an able-bodied control limb (CON). Data are presented as mean \pm SD for $n = 9$ (AMP and INT) and $n = 9$ (CON). Data in italics correspond to those variables where $n = 8$ due to participant withdrawal. Differences compared to AMP are denoted by * ($p < 0.05$) or ** ($p < 0.001$).

	AMP	INT	CON
Knee Extensor Kinetics			
Moment (Nm)	26.1 ± 13.3	65.4 ± 38.1 *	57.0 ± 13.7 *
Moment _{BM} (Nm.kg ⁻¹)	0.30 ± 0.14	0.75 ± 0.31 *	0.71 ± 0.24 *
Impulse (Nm·s)	1.14 ± 0.84	2.23 ± 1.21	1.75 ± 0.43
Impulse _{BM} (Nm·s.kg ⁻¹)	0.013 ± 0.009	0.025 ± 0.011	0.022 ± 0.008
Neural Drive			
Voluntary Activation (%)	50.6 ± 12.7	89.2 ± 5.75 **	90.4 ± 4.07 **
RMS EMG _{MVT} (% M _{max})	5.19 ± 1.20	9.10 ± 2.39 **	7.64 ± 1.47 *
Explosive RMS EMG ₀₋₁₀₀ (% M _{max})	5.38 ± 3.12	7.92 ± 3.66	7.00 ± 1.75
Evoked Twitch			
PT (Nm)	11.6 ± 6.00	30.8 ± 11.6 **	39.0 ± 11.9 **
Absolute RTD (Nm.s ⁻¹)	223 ± 171	650 ± 247 **	808 ± 243 **
Relative RTD (PT.s ⁻¹)	16.7 ± 3.23	20.4 ± 1.79 *	21.1 ± 4.60 *
Evoked Octet			
PT (Nm)	47.1 ± 31.2	94.5 ± 32.3 **	116 ± 28.0 **
Absolute RTD (Nm.s ⁻¹)	609 ± 387	1647 ± 541 **	1840 ± 365 **
Relative RTD (PT.s ⁻¹)	13.3 ± 1.62	17.7 ± 1.66 **	16.0 ± 1.43 *
Muscle Architecture			
Muscle Thickness (mm)	15.4 ± 5.19	26.3 ± 6.38 *	25.0 ± 3.34 *
Pennation Angle (°)	12.0 ± 1.66	13.9 ± 3.79	13.7 ± 1.46
Fascicle Length (mm)	73.8 ± 23.2	117 ± 50.8 **	96.5 ± 14.8

RTD, Rate of Torque Development; subscript BM, relative to body mass; subscript MVT, relative to Maximum Voluntary Torque; RMS EMG_{MVT}, root mean squared electromyography at MVT; RMS EMG₀₋₁₀₀, root mean squared electromyography from 0-100 ms of an explosive voluntary contraction; M_{max}, maximal M-wave; PT, peak torque.

699
700

Figure Headings

701 Figure 1. Static B-mode ultrasound image of the Vastus Lateralis (VL) and Vastus
702 Intermedius (VI) muscles at rest for the amputated (AMP) and intact (INT) limb of one ITTA,
703 and one control limb (CON). Architectural measures taken included pennation angle (Θ)
704 relative to the deep aponeurosis and extrapolated fascicle length, which were each
705 determined from three fascicles; and muscle thickness, measured between the superficial
706 and deep aponeuroses at three separate points (the centre and either end of each image –
707 indicated by numbered circles in the middle image). Significant reduction in amputated limb
708 VL muscle thickness is evident, while similarities in pennation angle in all three limbs, and
709 muscle thickness between INT and CON, can clearly be seen.

710 Figure 2. Maximal voluntary torque (MVT) and absolute peak rate of torque development
711 (RTD) recorded during respective maximal and explosive voluntary isometric knee
712 extensions, in both the amputated (AMP, light grey) and intact (INT, dark grey) limbs of
713 unilateral transtibial amputees ($n = 9$), and an able-bodied control group (CON, striped; $n =$
714 9). Data are presented as mean \pm SD absolute values (A,B) and relative to body mass (C) or
715 MVT (D). Differences compared to AMP are indicated by * ($p < 0.05$) or ** ($p < 0.001$).

716 Figure 3. Sagittal plane knee moments during the stance phase of walking for the amputated
717 (AMP, light grey line) and intact (INT, dark grey line) limbs of unilateral transtibial amputees,
718 and of an able-bodied control limb (CON, dashed line). INT and CON display substantial
719 overlap. Joint moment is expressed as internal moment. Positive and negative values
720 indicate knee extension and flexion moments, respectively. Data are presented as mean \pm
721 SD for $n = 8$ (AMP and INT) and $n = 9$ (CON).





